


## WARNING

KEEP OUT OF REACH OF CHILDREN

# TYPHOON



Active ingredient: Contains 15 g/litre tebuconazole and 15 g/litre boric acid in the form of an oil in water emulsion (EW)

A fungicide for the control of brown rot *Monolinia* spp on summerfruit.

Registered to:


Omnia Primaxa Limited  
PO Box 201041, Auckland Airport, Manukau 2150.  
Telephone: (09) 257 4030  
Facsimile: (09) 257 4031


In an emergency (All hours) Dial 0800 CHEMICAL 0800 243 622  
In a transport emergency Dial 111

Net:

Batch:

Date of Manufacture:

  
MARIE ZIVZELY  
Programme Manager  
(Operations)



## READ THE LABEL CAREFULLY BEFORE USING TYPHOON

**ACTIVE INGREDIENT:** Contains 15 g/litre tebuconazole and 15 g/litre boric acid in the form of an oil in water emulsion.

**HSNO CLASS: 6.3B, 6.8B, 6.9B, 9.1D.**

### TOXICITY:

- **HARMFUL:** May cause skin irritation
- **HARMFUL:** May cause reproductive /development damage from repeated oral exposure
- **HARMFUL:** May cause organ damage from repeated oral exposure at high doses

### ECOTOXICITY:

- **HARMFUL** to aquatic organisms.. Avoid contamination of any water supply with product or empty container

### PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- When mixing or applying Typhoon wear protective clothing such as impervious gloves and face protection to avoid contact with skin and eyes, and inhalation of spray mist.
- **Do not** eat, drink or smoke while using.
- Wash hands and face thoroughly after use and before eating or drinking.

### STORAGE:

- Store in original container, tightly closed and in a locked, dry, cool, well ventilated area away from foodstuffs, feeds, seeds, fertilisers, insecticides, or fungicides used for the protection of crops.
- Storage must be in accordance with the current version of NZS 8409 Management of Agrichemicals.

### SHELF LIFE:

- When stored appropriately, this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.



**SPILLAGE:**

- Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste container. Wash area with water and absorb with further inert material. Dispose of waste safely (such as a suitable landfill)

**DISPOSAL:**

- Triple rinse container and add residue to application container where possible. Crush and bury empty container in a suitable landfill.
- Avoid contamination of any water supply with chemical or empty container.

**FIRST AID:**

If swallowed DO NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin contact occurs, remove contaminated clothing and flush skin with running water. If splashed in eyes, wash out immediately with water.

**DIRECTIONS FOR USE:**


**Resistance Management:** Resistance to this fungicide and related DMI products could develop in some disease situations from repeated use. To minimise this risk use strictly in accordance with label instructions.

Apply a maximum of 3 spray applications of Typhoon, or any other DMI fungicide per season. Apply alternative non-DMI fungicides at other times.

**Compatibility:** Typhoon is compatible with most commonly used insecticides and fungicides. Do not use with alkaline material such as Bordeaux mixture and lime sulphur.

**Withholding period:** Summerfruit – 1 day

**Spray volume:** The recommendation given is for high volume spray applications to the point of run-off. For low volume applications make the necessary adjustment.



**Application:** For the control of brownrot on summerfruit

Crop	Disease	Dosage rate	Remarks
Summerfruit	Brownrot ( <i>Monolinia</i> spp)	665ml / 100 litres of water (Apply a minimum of 13.3 litres /ha)	<p>Apply as a high volume application to the point of run-off.</p> <p><b>Alternative 1.</b> Apply in a regular programme of sprays from pink to shuck fall. Thereafter apply alternative non-DMI fungicides through to harvest.</p> <p><b>Alternative 2.</b> Apply 2-3 sprays at 7-10 day intervals in the period prior to harvest. Use alternative non DMI sprays earlier in the season</p>

**CONDITIONS OF SALE:** The use of Typhoon being beyond the control of the suppliers, no warranty expressed or implied is given by Omnia Primaxa Ltd, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Omnia Primaxa Ltd accept no responsibility whatsoever resulting from the use of this product.

Registered pursuant to the ACVM Act 1997: P8087

See [www.nzfsa.govt.nz/acvm/](http://www.nzfsa.govt.nz/acvm/) for registration conditions.

Approved pursuant to the HSNO Act 1996, approval code: HSR007920

See [www.ernanz.govt.nz/search/registers.html](http://www.ernanz.govt.nz/search/registers.html) for controls applying to the substance.

