

7216  
3000


**KEEP OUT OF REACH OF CHILDREN**

# TACTIC

## FUNGICIDE

**ACTIVE INGREDIENT:** Contains 190 g/litre oleic acid as the potassium salt and 5g/litre salicylic acid in the form of a soluble concentrate.

A FUNGICIDE for the control of Botrytis on Grapes. Tactic can also be used as an adjuvant (wetting agent) for application of fungicides to a wide range of fruit, vegetable crops and ornamentals



**Registered to:**

Omnia Primaxa Ltd  
132 Pavillion Drive  
Airport Oaks  
Manukau  
2150

Telephone: (09) 257 4030  
Facsimile: (09) 257 4031


In an emergency (All hours)  
Call 0800 CHEMCALL 0800 243 622

In a transport emergency  
Call 111

**Net:**

**Batch:**

**Date of Manufacture:**



READ THE LABEL CAREFULLY BEFORE USING TACTIC  
ACTIVE INGREDIENT: Contains 190 g/litre oleic acid as the potassium salt and 5g/litre salicylic acid in the form of a soluble concentrate.

**HSNO CLASS:** 3.1C, 6.3B, 6.4A, 9.1D.

**WARNING:**

- FLAMMABLE LIQUID AND VAPOUR.

**TOXICITY:**

- HARMFUL: MAY CAUSE SKIN IRRITATION
- HARMFUL: MAY CAUSE EYE IRRITATION.

**ECOTOXICITY:**

- HARMFUL TO AQUATIC ORGANISMS. AVOID CONTAMINATION OF ANY WATER SUPPLY WITH CHEMICAL OR EMPTY CONTAINER.

**PRECAUTIONS:**

- KEEP OUT OF REACH OF CHILDREN.
- When mixing or applying, wear eye and face protection, waterproof gloves and overalls.
- Avoid contact with skin, eyes, clothes and inhalation of spray mist. Wash splashes off exposed skin immediately.
- Remove protective clothing and wash hands and face before meals and after work.
- Do not eat, drink or smoke while using

**DISPOSAL:**

- Container Disposal – triple rinse empty container and add rinsate to spray tank. Make sure the container is empty then burn in an appropriate incinerator, if circumstances such as wind direction permit. Otherwise crush or puncture and bury in a suitable landfill, or if appropriate, recycle.

**STORAGE:**


- Store in original container, tightly closed in a locked dry, cool, well ventilated area away from sources of ignition, foodstuffs, feeds, seeds, fertilisers, insecticides, or fungicides used for the protection of crops.
- Storage must be in accordance with the current version of NZS 8409 Management of Agrichemicals.
- Do not store with class 1, 2, 3, 2, 4 or 5 substances.

**SPILLAGE:**

- Wear appropriate protective clothing and prevent the chemical from entering waterways. Wash area with water and absorb with inert material. Dispose of waste safely in a suitable landfill.

**SHELF LIFE:**

- When stored appropriately Tactic should show no significant degradation for two years from the date of manufacture. Contact the supplier for further information about the use of any product that is older than this.



**FIRST AID:** If swallowed DO NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin contact occurs, remove contaminated clothing and flush skin with running water. If splashed in eyes, wash out immediately with water.

**GENERAL DIRECTIONS:**

Before opening, carefully read warning statement, precautions instructions, first aid instructions and directions for use.

**MIXING:**

**SHAKE OR STIR THE CONTENTS OF THE CONTAINER WELL BEFORE USE.**

Mix the required quantity of Tactic in the spray tank partly filled with water. Maintain agitation throughout the filling and spraying process.

**WITHHOLDING PERIODS:**

In all crops do not apply later than 14 days before harvest.

**DIRECTIONS FOR USE:**

Full spray coverage is essential.

The rates suggested are for high volume spraying to the point of run-off. For concentrate spraying, adjust dilution rate accordingly.

**For the control of Botrytis on grapes:** Apply 15 to 20 litres of Tactic in 500-1000 litres of water per ha. Apply Tactic as a preventative treatment when disease infection is expected in the period from before flowering to 14 days before harvest or at first signs of disease and then repeat at 14 day intervals depending on rainfall, inoculum levels and crop susceptibility.

Critical periods for botrytis control are over the flowering period, before bunch closure, veraison and in the period leading up to harvest.

For optimum results, apply a programme of sprays at 14 day intervals from early flowering to 14 days before harvest.

**For use as an adjuvant:** Apply 7.5 to 10 L of Tactic per Ha. For use with fungicides on a wide range of fruit, vegetable crops and ornamentals.

**COMPATIBILITY:** Tactic is compatible with Copper hydroxide (blackspot and downy mildew), Sulphur (powdery mildew), Captan (deadarm, blackspot and botrytis), Tolythuanid (500g/kg WG for botrytis, downy mildew and powdery mildew), Cyprodinil/Fludioxinil(375/250g/kg WG for botrytis), Delan (700 g/kg WG for blackspot and downy mildew), Potassium sulphate and the DMI / Triazole fungicides (powdery mildew).



**CONDITIONS OF SALE:** The use of Tactic being beyond the control of the suppliers, no warranty expressed or implied is given by Omnia Primaxa Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not, and Omnia Primaxa Ltd accept no responsibility whatsoever resulting from the use of this product.

Registered Pursuant to the ACVM Act 1997 No. P81112

See [www.nzfsa.govt.nz/acvm/](http://www.nzfsa.govt.nz/acvm/) for registration conditions.

Approved pursuant to the HSNO Act 1996, approval code HSR100026

See [www.ernanz.govt.nz/search/registers.html](http://www.ernanz.govt.nz/search/registers.html) for controls applying to the substance.

